

5TH

FILE 'HOME' ENTERED AT 14:23:21 ON 11 JUL 2003

L1 QUE (PATHOGEN OR BACTER#### OR VIRUS) (S) ((DEVELOPEMENT## OR DEVELOPMENT#  
#) (A) (DISORDER) OR AUTISM OR PDD)

L2 169 (PATHOGEN OR BACTER#### OR VIRUS) (S) ((DEVELOPEMENT## OR DEVELOPMENT##) (A) (DISORDER) OR AUTISM OR PDD)

L9 7 L8 AND (DIAGNOSE OR DETERMINE) (S) ((DEVELOPEMENT## OR DEVELOPMENT##) (A) DISORDER OR AUTISM OR PDD)

L10 18 L8 AND (DIAGNOS## OR DETERMIN#####) (S) ((DEVELOPEMENT## OR DEVELOPMENT##) (A) DISORDER OR AUTISM OR PDD)

L11 24 L7 AND (DIAGNOS## OR DETERMIN#####) (S) ((DEVELOPEMENT## OR DEVELOPMENT##) (A) DISORDER OR AUTISM OR PDD)

=> d his

(FILE 'HOME' ENTERED AT 14:23:21 ON 11 JUL 2003)

INDEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, AQUASCI, BIOBUSINESS, BIOCOMMERCE, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CANCERLIT, CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DDFB, DDFU, DGENE, DRUGB, DRUGLAUNCH, DRUGMONOG2, ...' ENTERED AT 14:23:46 ON 11 JUL 2003

SEA (PATHOGEN OR BACTER#### OR VIRUS) (S) ((DEVELOPEMENT## OR D

-----

12 FILE ADISCTI  
19 FILE ADISNEWS  
2 FILE AGRICOLA  
1 FILE AQUASCI  
1 FILE BIOBUSINESS  
1 FILE BIOCOMMERCE  
96 FILE BIOSIS  
169 FILE BIOTECHABS  
169 FILE BIOTECHDS  
24 FILE BIOTECHNO  
19 FILE CABA  
15 FILE CANCERLIT  
26 FILE CAPLUS  
1 FILE CONFSCI  
2 FILE DDFB  
3 FILE DDFU  
3212 FILE DGENE  
2 FILE DRUGB  
10 FILE DRUGU  
1 FILE EMBAL  
68 FILE EMBASE  
32 FILE ESBIODASE  
22\* FILE FEDRIP  
1 FILE FROSTI  
1 FILE FSTA  
1 FILE GENBANK  
1 FILE HEALSAFE  
10 FILE IFIPAT  
9 FILE JICST-EPLUS  
30 FILE LIFESCI  
36 FILE MEDLINE

2 FILE NIOSHTIC  
 2 FILE NTIS  
 34 FILE PASCAL  
 1 FILE PHARMAML  
 7 FILE PHIN  
 18 FILE PROMT  
 67 FILE SCISEARCH  
 25 FILE TOXCENTER  
 194 FILE USPATFULL  
 3 FILE USPAT2  
 1 FILE VETU  
 172 FILE WPIDS  
 172 FILE WPINDEX

L1 QUE (PATHOGEN OR BACTER#### OR VIRUS) (S) ((DEVELOPEMENT## OR D  
 -----

FILE 'BIOTECHABS' ENTERED AT 14:28:24 ON 11 JUL 2003

L2 169 S L1  
 L3 169 DUP REM L2 (0 DUPLICATES REMOVED)  
 L4 169 S L3  
 L5 14 S L3 NOT PY>2000

FILE 'MEDLINE, CAPLUS, BIOSIS, BIOTECHNO, EMBASE, SCISEARCH' ENTERED AT  
 14:31:48 ON 11 JUL 2003

L6 317 S L1  
 L7 152 DUP REM L6 (165 DUPLICATES REMOVED)  
 L8 89 S L7 NOT PY>2000  
 L9 7 S L8 AND (DIAGNOSE OR DETERMINE) (S) ((DEVELOPEMENT## OR DEVEL  
 L10 18 S L8 AND (DIAGNOS## OR DETERMIN#####) (S) ((DEVELOPEMENT## OR  
 L11 24 S L7 AND (DIAGNOS## OR DETERMIN#####) (S) ((DEVELOPEMENT## OR  
 L12 6 S L11 NOT L10

=>

L10 ANSWER 2 OF 18 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 AN 2000:457217 BIOSIS  
 DN PREV200000457217  
 TI Enterocolitis in children with developmental disorders.  
 AU Wakefield, A. J. (1); Anthony, A.; Murch, S. H.; Thomson, M.; Montgomery, S. M.; Davies, S.; O'Leary, J. J.; Berelowitz, M.; Walker-Smith, J. A.  
 CS (1) Inflammatory Bowel Disease Study Group, Department of Medicine, Royal Free and University College Medical School, Hampstead, Royal Free Campus, London, NW3 2QG UK  
 SO American Journal of Gastroenterology, (September, 2000) Vol. 95, No. 9, pp. 2285-2295. print.  
 ISSN: 0002-9270.  
 DT Article  
 LA English  
 SL English  
 AB OBJECTIVE: Intestinal pathology, i.e., ileocolonic lymphoid nodular hyperplasia (LNH) and mucosal inflammation, has been described in children with **developmental disorders**. This study describes some of the endoscopic and pathological characteristics in a group of children with **developmental disorders** (affected children) that are associated with behavioral regression and bowel symptoms, and compares them with pediatric controls. METHODS: Ileocolonoscopy and biopsy were performed on 60 affected children (median age 6 yr, range 3-16; 53 male). Developmental **diagnoses** were **autism** (50 patients), Asperger's syndrome (five), disintegrative disorder (two), attention deficit hyperactivity disorder (ADHD) (one), schizophrenia (one), and dyslexia (one). Severity of ileal LNH was graded (0-3) in both affected children and 37 developmentally normal controls (median age 11 yr, range 2-13 yr) who were investigated for possible inflammatory bowel disease (IBD). Tissue sections were reviewed by three pathologists and scored on a standard proforma. Data were compared with ileocolonic biopsies from 22 histologically normal children (controls) and 20 children with ulcerative colitis (UC), scored in an identical manner. Gut **pathogens** were sought routinely. RESULTS: Ileal LNH was present in 54 of 58 (93%) affected children and in five of 35 (14.3%) controls ( $p < 0.001$ ). Colonic LNH was present in 18 of 60 (30%) affected children and in two of 37 (5.4%) controls ( $p < 0.01$ ). Histologically, reactive follicular hyperplasia was present in 46 of 52 (88.5%) ileal biopsies from affected children and in four of 14 (29%) with UC, but not in non-IBD controls ( $p < 0.01$ ). Active ileitis was present in four of 51 (8%) affected children but not in controls. Chronic colitis was identified in 53 of 60 (88%) affected children compared with one of 22 (4.5%) controls and in 20 of 20 (100%) with UC. Scores of frequency and severity of inflammation were significantly greater in both affected children and those with UC, compared with controls ( $p < 0.001$ ). CONCLUSIONS: A new variant of inflammatory bowel disease is present in this group of children with **developmental disorders**.

L10 ANSWER 5 OF 18 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 AN 2000:39700 BIOSIS  
 DN PREV200000039700  
 TI Gastrointestinal abnormalities in children with autistic disorder.  
 AU Horvath, Karoly (1); Papadimitriou, John C.; Rabszty, Anna; Drachenberg, Cynthia; Tyson Tildon, J.  
 CS (1) Department of Pediatrics, 22 S Greene St, N5W70, Baltimore, MD, 21201-1595 USA  
 SO Journal of Pediatrics, (Nov., 1999) Vol. 135, No. 5, pp. 559-563.  
 ISSN: 0022-3476.  
 DT Article

LA English  
 SL English  
 AB Objectives: Our aim was to evaluate the structure and function of the upper gastrointestinal tract in a group of patients with **autism** who had gastrointestinal symptoms. Study design: Thirty-six children (age: 5.7 +- 2 years, mean +- SD) with autistic disorder underwent upper gastrointestinal endoscopy with biopsies, intestinal and pancreatic enzyme analyses, and **bacterial** and fungal cultures. The most frequent gastrointestinal complaints were chronic diarrhea, gaseousness, and abdominal discomfort and distension. Results: Histologic examination in these 36 children revealed grade I or II reflux esophagitis in 25 (69.4%), chronic gastritis in 15, and chronic duodenitis in 24. The number of Paneth's cells in the duodenal crypts was significantly elevated in autistic children compared with non-autistic control subjects. Low intestinal carbohydrate digestive enzyme activity was reported in 21 children (58.3%), although there was no abnormality found in pancreatic function. Seventy-five percent of the autistic children (27/36) had an increased pancreatico-biliary fluid output after intravenous secretin administration. Nineteen of the 21 patients with diarrhea had significantly higher fluid output than those without diarrhea. Conclusions: Unrecognized gastrointestinal disorders, especially reflux esophagitis and disaccharide malabsorption, may contribute to the behavioral problems of the non-verbal autistic patients. The observed increase in pancreatico-biliary secretion after secretin infusion suggests an upregulation of secretin receptors in the pancreas and liver. Further studies are required to **determine** the possible association between the brain and gastrointestinal dysfunctions in children with autistic disorder.

L10 ANSWER 7 OF 18 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 AN 1998:258432 BIOSIS  
 DN PREV199800258432  
 TI Bovine virus diarrhoea. Review article.  
 AU Kecskemeti, S. (1); Kiss, I.  
 CS (1) Debreceni Allat-egeszsegugyi Intezet, Bornemissza u. 3-5, H-4031 Debrecen Hungary  
 SO Magyar Allatorvosok Lapja, (March, 1998) Vol. 120, No. 3, pp. 143-151. ISSN: 0025-004X.  
 DT Article  
 LA Hungarian  
 SL Hungarian; English  
 AB The infection caused by bovine diarrhoea **virus** (BVDV) generally causes mild clinical symptoms in immunocompetent cattle. A severe disease can be observed mainly in young animals. The infection of pregnant animals is especially important because the **virus** may invade the fetus. Due to an infection caused by a non-cytopathogenic BVDV strain persistently infected, immunotolerant animals may be born which are consistently viraemic and can shed the **virus** during the whole lifetime. Such an infection can not be caused by a cytopathogenic biotype. **Developmental disorders** of central nervous system and eyes may develop in case of an infection between the 90th and 150th days of pregnancy. The fetus is able to produce antibodies against the **virus** during the second half of pregnancy. Venereal infections are also important. Bulls - in case of an acute infection with lower titres and for a shorter period, however in case of persistent infections persistently and with a higher titre - shed the **virus**. The **virus** in the semen may cause seroconversion, return to heat, embryonic disorders, etc. in the infected animals and the animals born can be persistently infected. Mucosal disease (MD) may develop in animals persistently infected with a non-cytopathogenic BVDV during the

fetal life - between the 40th and 120th days - when the animals are super infected with a cytopathogenic BVDV later on. When the super infecting **virus** has a homologous antigenic structure a disease develops with low morbidity but high mortality. When the antigenic structure of the super infecting **virus** is partly heterologous, due to the antibodies produced against it the super infecting **virus** disappear from the blood. MD develops weeks or months after the super infection. MD does not develop after a super infection with homologous or heterologous cytopathogenic BVDV, even antibodies are produced against the heterologous **virus**. **Diagnosis** of the diseases caused by BVDV is based on the clinical symptoms, pathological and histological alterations and results of laboratory investigations. Demonstration of the **virus**, **virus** antigen or nucleic acid are used for the laboratory **diagnosis** of BVDV.

L10 ANSWER 8 OF 18 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 AN 1995:228097 BIOSIS  
 DN PREV199598242397  
 TI Prevalence of hepatitis B virus markers in several institutions for the mentally handicapped in the autonomous community of Madrid.  
 AU Carrascosa, Diana; De La Vega Ramirez, M.; Casado, Angela; Je La Torre, M. Roserio; Lopezfernandez, M. Encarnacion; Saez, Julia  
 CS Departamento de Fisiopatologia y Genetica Molecular Humana, Centro di Investigaciones Biologicas, c. Velazquez 144, 28006-Madrid Spain  
 SO American Journal of Human Biology, (1995) Vol. 7, No. 2, pp. 217-222. ISSN: 1042-0533.  
 DT Article  
 LA English  
 AB In order to **determine** the prevalence of hepatitis B **virus** (HBV) markers, 400 patients were studied: 134 residents of an institution (RI) for the mentally retarded and 266 under non-residential care (NRC). In the residential institutions, all markers were absent in 69 (65.7%) of 105 patients with Down syndrome and 20 (69.0%) of 29 clients without Down syndrome. In the NRC clients, 167 (85.6%) of 195 patients with Down syndrome and 65 (91.5%) of 71 clients with other mental defects (psychologically and physically handicapped, **autism**) had negative tests for HBV markers. The prevalence of the hepatitis B surface antigen (HBsAg) was higher in institutionalized mentally retarded (RI) and older patients (21+ years). Examination of 195 Down patients revealed a higher frequency (1.4 times) of surface antigen carriers as strictly matched nonDown syndrome cases (point prevalences 14.2% and 10.3%, respectively). The higher prevalence in affected cases appears to be primarily associated with a longer persistence of antigenemia. Results related to the sex of the patients were less clear. Neither affected nor non-affected patients showed significant differences in prevalence among males and females.

L10 ANSWER 10 OF 18 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 AN 1993:505907 BIOSIS  
 DN PREV199396129914  
 TI Infantile autism in Nordland County: Prevalence and etiology.  
 AU Herder, Gyro Aas  
 CS Barneavdelingen, Nordland Sentralsykehus, 8017 Bodo Norway  
 SO Tidsskrift for den Norske Laegeforening, (1993) Vol. 113, No. 18, pp. 2247-2249. ISSN: 0029-2001.  
 DT Article  
 LA Norwegian  
 SL Norwegian; English  
 AB Few epidemiological studies of infantile **autism** have been

conducted in Norway. A prevalence of four per 100 000 children was found in Nord-Trøndelag. Studies from Sweden, England and Japan show prevalence figures of 2-13 per 10 000. Causal factors of **autism** are complicated, but many studies in recent years indicate that both genetic and neurological factors are important. The aim of this study was to establish the prevalence of infantile **autism** in Nordland County look for casual factors and disabilities among the children. Together with the **Autism Team** we found 28 persons, born between 1975 and 1991, who lived in the county in 1992. This gives a prevalence of 5.5 per 10 000. 21 were boys, seven girls. 26 were mentally retarded, 13 had no verbal speech. Nine had epilepsy. 13 children were **diagnosed** before the age of four. Different diseases and impairments associated with **autism** were discovered in all 19 children. In the case of eight children we found associated disease. When a **diagnosis** of infantile **autism** is made, the child should undergo a thorough medical and neurological examination. This should include CT brain scan, EEG, chromosome analysis, and screening for metabolic diseases and intrauterine **virus** infections. Sometimes examinations of liquor and auditory brain-stem responses are indicated.

- L10 ANSWER 13 OF 18 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 AN 1984:203295 BIOSIS  
 DN BA77:36279  
 TI AUTISM IN A CHILD WITH CONGENITAL CYTOMEGALOVIRUS INFECTION.  
 AU MARKOWITZ P I  
 CS DEP. PEDIATR., MED. COLL. VIRGINIA, BOX 487, RICHMOND, VIRGINIA 23298.  
 SO J AUTISM DEV DISORD, (1983) 13 (3), 249-254.  
 CODEN: JADDDQ.  
 FS BA; OLD  
 LA English  
 AB A case is reported of early infantile **autism** associated with a congenital cytomegalovirus infection. The **diagnosis** of **autism** is based on the child's failure to develop good interpersonal relationships, poor eye contact, delayed and deviant use of language and her rote and nonthematic use of objects and playthings. Resistance to change and self-stimulatory behavior were also present. Onset was before 2 yr of age. Congenital cytomegalovirus infection was suggested by the presence of an antibody response to the **virus**, culture of the **virus** from the urine, sensorineural hearing loss and inflammatory damage to the retina of the eye. Although over time improvement was noted, at last examination at the age of 5 yr her behavior is still markedly deviant. This and other reported cases suggest that congenital viral infection may be an important cause of infantile **autism**. It is hypothesized that an ability of the agent to establish chronic infection may predispose to behavioral aberration.
- L10 ANSWER 15 OF 18 EMBASE COPYRIGHT 2003 ELSEVIER SCI. B.V.  
 AN 1998263515 EMBASE  
 TI Possible immunogenetic basis for autism.  
 AU Burger R.A.; Warren R.P.  
 CS R.A. Burger, Utah State University, Logan, UT 84322-6895, United States.  
 Roger@cpd2.usu.edu  
 SO Mental Retardation and Developmental Disabilities Research Reviews, (1998) 4/2 (137-141).  
 Refs: 28  
 ISSN: 1080-4013 CODEN: MRDRFI  
 CY United States  
 DT Journal; General Review  
 FS 022 Human Genetics  
 026 Immunology, Serology and Transplantation

032      Psychiatry

LA    English

SL    English

AB    **Autism** results from several different etiologies or combination of pathological mechanisms. Mounting evidence indicates that immune dysfunction along with an environmental **pathogen** may be factors contributing to the development of some cases of **autism**. One of the immune deficiencies observed in **autism** is abnormal T-cell mediated immunity. Another is altered levels of certain classes of antibodies (immunoglobulins), including decreased levels of immunoglobulin A and deficient complement activity, based on the inheritance of a null allele of the C4B gene. In addition to the C4B gene, other genes on chromosome 6 also appear to be associated with **autism**. In the developing child, genetically **determined** immune deficiencies might increase the risk for **autism** in two ways: 1) A **pathogen** or its toxins might damage the brain, 2) the **pathogen** might trigger an autoimmune mechanism that would interfere with brain functioning. In the mother, immune deficiency might allow a **pathogen** to persist in utero, damaging the fetal brain directly or triggering a maternal immune response that creates pathogenesis in the fetal brain.

L12 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2003 ACS  
 AN 2002:488136 CAPLUS  
 DN 137:30245  
 TI Methods for diagnosing pervasive development disorders, dysautonomia and  
 other neurological conditions  
 IN Fallon, Joan M.  
 PA USA  
 SO U.S. Pat. Appl. Publ., 9 pp.  
 CODEN: USXXCO  
 DT Patent  
 LA English  
 FAN.CNT 1

|      | PATENT NO.      | KIND | DATE     | APPLICATION NO. | DATE     |
|------|-----------------|------|----------|-----------------|----------|
|      | -----           | ---- | -----    | -----           | -----    |
| PI   | US 2002081628   | A1   | 20020627 | US 2001-990909  | 20011116 |
| PRAI | US 2000-249239P | P    | 20001116 |                 |          |

AB Methods for aiding in the **diagnosis** of disorders including, but  
 not limited to, **PDDs** (Pervasive Development  
**Disorders**), Dysautonomic disorders, Parkinson's disease and SIDS  
 (Sudden Infant Death Syndrome). In one aspect, a **diagnosis**  
 method comprises analyzing a stool sample of an individual for the  
 presence of a biol. marker (or marker compd.) comprising one or more  
**pathogens**, which provides an indication of whether the individual  
 has, or can develop, a disorder including, but not limited to, a  
**PDD**, Dysautonomia, Parkinsons disease and SIDS. Preferably, the  
 presence of one or more pathogens is detd. using a stool immunoassay to  
 det. the presence of antigens in a stool sample, wherein such antigens are  
 assocd. with one or more pathogens including, but not limited to, Giardia,  
 Cryptosporidium, E. histolytica, C. difficile, Adenovirus, Rotavirus or H.  
 pylori.



L13 10421 (DISEASE OR DISORDER) (S) (PLURAL### OR MULTIPL##### OR SEVERAL)  
 (3A) (PATHOGEN OR BACTER### OR VIRUS## OR INFECT####)  
 L14 2799 L13 AND (DIAGNOS### OR DETERMIN#####) (S) (DISEASE OR DISORDER)  
 L15 2700 L13 AND ((DISEASE OR DISORDER) (S) (PLURAL### OR MULTIPL#####  
 OR SEVERAL) (3A) (PATHOGEN OR BACTER### OR VIRUS## OR INFECT####  
 ) ) (P) ((DIAGNOS### OR DETERMIN#####) (S) (DISEASE OR DISORDER)  
 )  
 L13 10421 S (DISEASE OR DISORDER) (S) (PLURAL### OR MULTIPL##### OR SEVER  
 L14 2799 S L13 AND (DIAGNOS### OR DETERMIN#####) (S) (DISEASE OR DISORDE  
 L15 2700 S L13 AND ((DISEASE OR DISORDER) (S) (PLURAL### OR MULTIPL#####  
 L16 1333 DUP REM L15 (1367 DUPLICATES REMOVED)  
 L17 1025 S L16 NOT PY>2000  
 L18 10 S L17 AND (HELICOBACTER OR PYLORI)

(DIAGNOS### OR DETERMIN#####) (S) (DISEASE OR DISORDER)

L18 ANSWER 3 OF 10 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

AN 2000:169707 BIOSIS

DN PREV200000169707

TI Patients with acute myocardial infarction in Northern Italy are often infected by **Helicobacter pylori**.

AU Pellicano, R.; Parravicini, P. P.; Bigi, R.; La Rovere, M. T.; Baduini, G.; Gandolfo, N.; Casaccia, M.; Reforzo, F.; Santoriello, L.; Aruta, E.; Marengo, G.; Arena, V.; Bazzoli, F.; Rizzetto, M.; Ponzetto, A. (1)

CS (1) UOA Gastroenterology, Molinette Hospital, Corso Bramante 88, 10126, Turin Italy

SO Panminerva Medica., (Dec., 1999) Vol. 41, No. 4, pp. 279-282.  
ISSN: 0031-0808.

DT Article

LA English

SL English

AB Background: The classical risk factors for acute myocardial infarction (AMI) fail to explain all the epidemiological variations of the **disease**. Among the new risk factors recently reported, **several infectious** agents appear to increase the risk of AMI. In particular, acute and chronic respiratory **diseases** due to Chlamydia pneumoniae, and **Helicobacter pylori** (H. **pylori**) infection seem to be strongly involved. The aim of this work is to **determine** the prevalence of H. **pylori** infection in a group of male patients with AMI, in a case-control study, where a group of blood donors matched for sex and age served as control. We searched for the classical risk factors in all patients. Methods: We studied 212 consecutive male patients, aged 40-65 years, admitted for AMI at the Coronary Care Units at Hospitals in three towns of Northern Italy. H. **pylori** infection was assessed by the highly specific and sensitive 13C-urea breath test and by presence of antibodies (IgG) against H. **pylori** in circulation. Volunteer blood donors attending our Hospital Blood Bank served as controls. Among the patients we investigated the presence of hypertension, cholesterol and glucose levels in serum, fibrinogen in plasma and the smoking habit. Results: H. **pylori** infection was present in 187/212 (88%) of the patients and in 183/310 (59%) of the control population ( $p < 0.0001$ ). Classical risk factors for AMI did not differ among patients with and without H. **pylori** infection. Conclusion: Patients admitted to the Coronary Care Unit for acute myocardial infarction had a notably higher prevalence of H. **pylori** infection than the general population. The classical risk factors for coronary **disease** were equally present in all patients with AMI irrespective of H. **pylori** status.

# WEST Search History

DATE: Friday, July 11, 2003

## Set Name Query side by side

## Hit Count Set Name result set

*DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=OR*

|     |   |      |     |
|-----|---|------|-----|
| L16 | L14 and (pathogen or virus or bacter\$4 or helicobacter or pylori) with autism  | 0    | L16 |
| L15 | L14 and (pathogen or virus or bacter\$4 or helicobacter or pylori) with (development\$3 adj disorder or autism or PDD)  | 1    | L15 |
| L14 | L13 not l4  | 155  | L14 |
| L13 | L12 and l3  | 167  | L13 |
| L12 | L11 or l8   | 580  | L12 |
| L11 | L10 and l5  | 580  | L11 |
| L10 | (pathogen or virus or bacter\$4 or helicobacter or pylori) and (development\$3 adj disorder or autism or PDD)   | 1449 | L10 |
| L9  | L7 and developmental adj disorder   | 0    | L9  |
| L8  | L7 and developmental adj disorder   | 150  | L8  |
| L7  | L6 not l4   | 155  | L7  |
| L6  | l3 and L5   | 167  | L6  |
| L5  | (pathogen or virus or bacter\$4 or helicobacter or pylori) and (development\$3 adj disorder or autism or PDD)   | 580  | L5  |
| L4  | L3 and l2   | 12   | L4  |
| L3  | (detect\$5 or determin\$5) with (pathogen or virus or bacter\$4 or helicobacter or pylori) and (determin\$5 or diagnos\$3) with (development adj disorder or autism or PDD) | 167  | L3  |
| L2  | (pathogen or virus or bacter\$4 or helicobacter or pylori) same (development adj disorder or autism or PDD)   | 106  | L2  |
| L1  | (pathogen or virus or bacter\$4 or helicobacter or pylori) and (development adj disorder or autism or PDD)  | 580  | L1  |

END OF SEARCH HISTORY

## WEST Search History

DATE: Friday, July 11, 2003

| <u>Set</u><br><u>Name</u><br>side by<br>side | <u>Query</u>  | <u>Hit</u><br><u>Count</u> | <u>Set</u><br><u>Name</u><br>result<br>set |
|--|---|----------------------------|--|
|  | <i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=OR</i>   |                            |  |
| L10  | (disease or disorder) with (plural\$4 or multipl\$4) adj3 pathogen\$2   | 44                         | L10  |
| L9   | L8 and l3   | 47                         | L9   |
| L8   | (plural\$4 or multipl\$5 or several) adj3 (antigen\$2 or pathogen\$2 or infect\$5)<br>same (diagnos\$4 or determin\$6) with (disease or disorder) | 109                        | L8   |
| L7   | (plural\$4 or multipl\$5 or several) adj3 (antigen\$2 or pathogen\$2 or infect\$5)<br>same (diagnos\$4 or determin\$6 or disease or disorder)     | 1349                       | L7   |
| L6   | l4 and l3   | 65                         | L6   |
| L5   | l4 and l3L4   | 0                          | L5   |
| L4   | l1 and (plural\$4 or multipl\$5 or several) adj3 (antigen\$2 or pathogen\$2 or<br>infect\$5)  | 145                        | L4   |
| L3   | L1 and @pd<20001116   | 170                        | L3   |
| L2   | L1 and @ad<20001116   | 254                        | L2   |
| L1   | (diagnos\$3 or determin\$6) with (disease or disorder) same (plural\$4 or<br>multipl\$5 or several) with (antigen\$2 or pathogen\$2 or infect\$5) | 352                        | L1   |

END OF SEARCH HISTORY

STIC-ILL

---

From: STIC-Biotech/ChemLib  
Sent: Friday, July 11, 2003 12:07 PM  
To: STIC-ILL  
Subject: FW: article request

Message to you.

-----Original Message-----

From: Lucas, Zacharia  
Sent: Friday, July 11, 2003 11:55 AM  
To: STIC-Biotech/ChemLib  
Subject: article request

Examiner# : 79253 Zachariah Lucas  
Art Unit : 1648  
Phone Number: 308-4240  
Date: 7-11-2003  
Serial Number: 09/990909  
MailBox & Bldg/Room Location: 8e12/8d16  
Results Format Preferred (circle): Paper

Could you please send me a copy of the following article(s).

ARchives of Disease in Childhood, June 2001, 84(6): 525, articles by Richardson, and Murphy

Thank you,  
Zac Lucas

## LETTERS TO THE EDITOR

The discovery that *Helicobacter pylori* is the prime cause of peptic ulcer disease, is one of the most important advances in medicine in the 20th century. Subsequently, its importance in the causation of gastric cancer has been recognised. It is a rare cause of gastric lymphoma. Despite its significance as a pathogen, this organism colonises the gastric mucosa in up to 50 percent of the world's population. Not surprisingly research interest is intense. There has been much speculation (though little proof) that it might have a role in various other gastrointestinal and non-gastrointestinal disorders, including failure to thrive in infancy, short stature, anaemia, and even cardiovascular disease. Now a link has been proposed between *H pylori* and sudden infant death syndrome (SIDS). Recently, Kerr *et al* examined gastric, tracheal, and pulmonary tissue, looking for evidence of *H pylori* in SIDS victims and controls. Based on polymerase chain reaction (PCR) techniques, they reported a highly significant association between SIDS and the presence of two *H pylori* genes (UreC, *cagA*) in these tissues. Not surprisingly, this reported association has evoked a lively correspondence. Important questions have been raised regarding both methodology and interpretation.

M STEPHEN MURPHY  
Associate Editor

1 Kerr JR, Al-Khattaf A, Barton AJ, *et al*. An association between sudden infant death syndrome (SIDS) and *Helicobacter pylori* infection. *Arch Dis Child* 2000;83:429-34.

### Association between SIDS and *H pylori* infection

**EDITOR.**—The article in the November issue of the *Archives* on the association between sudden infant death syndrome (SIDS) and *H pylori* infection is confusing.<sup>1</sup> I am very familiar with *H pylori* colonisation in gastric biopsies in children and its association with gastritis, peptic ulcer, and gastric cancer. However, the implication that the organism can cause an unexpected infant death that is, SIDS, is shocking!

Unexplained infant deaths (SIDS) are "fertile soil" for speculators that apply new technology polymerase chain reaction (PCR) to uncover new associations. Unfortunately, these observations are not based on an infrastructure of knowledge of the causes of infant mortality. Caution needs to be exercised when applying PCR technology to postmortem tissue and "discovering" an answer. The possibility of contamination is real, and in addition infants can die with something, and not of it.

I would value a response from Drs Fleming, Blair, Bacon, and Berry who co-authored the CESDI study of SUDI.

RAI LPH A FRANCIOSI  
Paediatric Pathologist,  
Children's Hospital and Medical College of Wisconsin,  
Wiscasset, WI, USA  
ralf@mc.wisc.edu

1 Kerr JR, Al-Khattaf A, Barton AJ, *et al*. An association between sudden infant death syndrome (SIDS) and *Helicobacter pylori* infection. *Arch Dis Child* 2000;83:429-34.

### Ammonia—not the culprit

**EDITOR.**—We were interested to read the article by Kerr *et al* on the SIDS problem. With regard to the interesting results we would like to point out some related findings. As pointed out by Kerr *et al*, *H pylori* is abundant in less advantageous parts of society where smoking is often frequent, and sometimes where SIDS occurs. The fact that smoking is often inversely related to the ability of *H pylori* to colonise and to be transmitted from mother to child<sup>2</sup> might indicate that it is sensitive to smoke itself, or products generated after smoke inhalation. It is interesting to note that endogenous products of smoke, like nitrate and nitrite, often inhibit bacterial growth.<sup>3</sup>

Furthermore, we have previously shown that total breakdown of all ingested urea takes place in all normal infants without causing problems of ammonia intoxication.<sup>4</sup> This is in contrast to SIDS victims, most of whom have unmetabolised urea in their faeces.<sup>5</sup> Due to these related circumstances it may seem a little adventurous to suggest that ammonia produced by *H pylori* could cause death in SIDS.

LARS WIKLUND  
GUNNAR RÖNQVIST  
MARY GEORGE  
Department of Anaesthesia,  
Uppsala University Hospital,  
Uppsala, Sweden  
Lars.Wiklund@anest.uu.se, lars.wiklund@anest.uu.se

- 1 Urener H, Bode G, Adler G, *et al*. Does maternal smoking hinder mother-child transmission of *Helicobacter pylori* infection? *Epidemiology* 2000;11:71-5.
- 2 Goreski J, Zafriou OC, Hollcher TC. Steady-state nitric oxide concentrations during denitrification. *J Biol Chem* 1990;265:1535-8.
- 3 George M, Nord KE, Rönquist G, *et al*. Faecal microflora and urease activity during the first six months of infancy. *Ups J Med Sci* 1996;101:231-50.
- 4 Wiklund L, George M, Nord KE, *et al*. Sudden infant death syndrome and nitrogen metabolism: further development of a hypothesis. *Eur J Clin Invest* 1998;28:958-65.

### Control your controls and conclusions

**EDITOR.**—In a retrospective study, Kerr and coworkers investigated formalin fixed, paraffin embedded tissues (stomach, trachea, and lung) of 32 infants who had died of SIDS, and eight control cases, with nested polymerase chain reaction (PCR) and ELISA of the amplicons. A child was considered as infected with *H pylori* if the optical density of the ELISA was above the mean value plus 2 SD obtained in the tissue of control infants. The authors found that 28 of the 32 SIDS cases, but only one of the eight control cases fulfilled these criteria. They conclude from their results that *H pylori* infection may play a causative role in SIDS. We have serious doubts about their results and conclusions.

The control group was extremely small in size and we would expect most, if not all, of

these eight infants to have received one or more antibiotics in high doses intravenously over several days before death, as the causes of death were bacterial meningitis, septicæmia, pneumonia, necrotising enterocolitis, ileal perforation, and prematurity. In contrast, few if any of the SIDS victims would have received intravenous antibiotics. Therefore, if control children had been colonised with *H pylori*, the bacteria may have been suppressed. These eight infants are certainly not appropriate controls for this kind of study.

Nested PCR is a very sensitive method with a high risk of false positive results caused by contamination. The applied ELISA is yet another amplifying method which also increases the risks of unspecific binding. Although the authors stated that they tried to minimise contamination, no precautions have been performed at the time of autopsy and preservation of the tissue due to the retrospective character of the study. Because of the low specificity of the methods used, it is mandatory to prove the identity of the PCR amplicons as *H pylori* specific by sequencing the products. Such confirmation is not reported in the paper. To show the specificity of their method, the authors could have also performed analyses on control tissues—for example, brain, which are unlikely to be *H pylori* infected even when other tissues were assessed as "positive".

The fact that *H pylori* was not shown in the stomach, trachea, or lung by histology in any of the children must raise major concerns that the applied methods were not specific. Other methods for detection of *H pylori* infection like fluorescence in situ hybridisation (FISH) have not been applied.<sup>6</sup> The authors do not report whether any of the children had histological signs of acute or chronic gastritis, which is found even in young children with *H pylori* infection.<sup>7</sup> If the bacterial load was so small that neither the bacteria nor the associated inflammation could be detected by histology, it seems questionable that metabolic products produced by *H pylori* (for example, ammonia) may play a causative role as a cause of SIDS as suggested by the authors.

Finally, the authors mention that both *H pylori* infection and SIDS are more common in poor socioeconomic populations but fail to provide any information on the ethnic and socioeconomic background of their cases and control infants. From many epidemiologic studies and our own experience, it seems extremely unlikely that 28 of 32 infants (87%) under 28 weeks of age are infected by *H pylori* in a country such as the UK, unless these children are from immigrant groups. We are, for example, following a cohort of German children from birth with regular testing for *H pylori* infection by two non-invasive tests: the detection of *H pylori* antigen in stool (HpSA, Meridian Diagnostics, Cincinnati, USA) and the "C-urea breath test corrected for estimated individual CO<sub>2</sub> production rate".<sup>8</sup> Although a quarter of the children have at least one *H pylori* infected parent (positive serology and/or a positive "C-urea breath test") only 1.5% of the children have positive tests during the first three years of age.

On publication, this paper was widely reported by the media, a process actively assisted by the authors. This is likely to result in considerable anxiety among young parents and pregnant women, feelings of guilt in parents of SIDS children and unjustified *H pylori*